



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

241. Proposed by CLIFFORD N. MILLS, Brookings, S. Dak.

If $a^2 + b^2 = c^2$, where a , b , and c are integers, then prove that abc will be a multiple of 60.

In the next issue we shall reprint all unsolved problems in Number Theory published since January, 1913. They are numbers 191, 192, 196, 198, 201, 202, 205, 208, 209, 211, 214, 217, 219, 221, 222, 223. Please have these in mind. EDITORS.

SOLUTIONS OF PROBLEMS.**ALGEBRA.****433. Proposed by B. J. BROWN, Student at Drury College.**

Prove that, if all the quantities, a , b , etc., are real, then all the roots of the equations

$$\begin{vmatrix} a-x & h \\ h & b-x \end{vmatrix} = 0, \quad \begin{vmatrix} a-x & h & g \\ h & b-x & f \\ g & f & c-x \end{vmatrix} = 0$$

are real; and generalize the proposition.

SOLUTION BY WM. E. HEAL, Washington, D. C.

The first equation may be written

$$\left[x - \left(\frac{a+b}{2} \right) \right]^2 = \frac{(a-b)^2}{4} + h^2.$$

Since the second member is the sum of two squares and so can never become negative, if a , b , and h are real, it follows that both roots are real.

The second equation is proved, in Salmon's *Modern Higher Algebra*, 4th edition, page 28, to have its roots all real.

The general equation, of which the above are special cases, is shown on page 48 of the same work to have all its roots real.

Thus also referred to by A. M. HARDING.

442. Proposed by CLIFFORD N. MILLS, Brookings, S. Dak.

Show that the sum of n terms of the series $1/2 - 1/3 + 1/4 - 1/6 + 1/8 - 1/12 + \dots$ is $1/3[1 - (1/2)^{n/2}]$ when n is even, and $1/3[1 + 2\sqrt{2}(1/2)^{(n/2)+1}]$ when n is odd.

SOLUTION BY IRBY C. NICHOLS, Chicago, Ill.

(1) *When n is even.* Grouping the terms successively by twos, we have a series of $n/2$ terms from which the factor $(1/2 - 1/3)$ may be removed, thus,

$$(1/2 - 1/3) \left[\frac{1}{2^0} + \frac{1}{2^1} + \frac{1}{2^2} + \dots + \frac{1}{2^{n/2-1}} \right].$$

The series in brackets is geometrical, and we have the sum

$$(1/2 - 1/3) \left[\frac{1 - 1/2 \cdot \frac{1}{2^{n/2-1}}}{1/2} \right] = 1/3[1 - (1/2)^{n/2}].$$

(2) *When n is odd.* Then the sum of $n+1$ terms can be written by (1), using $n+1$ for n . If now we add the $(n+1)$ th term to this sum, we shall have the sum of n terms, since the $(n+1)$ th term is negative. This gives

$$\begin{aligned} S_n &= 1/3[1 - (1/2)^{(n+1)/2}] + (1/2)^{[(n+1)/2]-1} \\ &= 1/3[1 - 2^{1/2}(1/2)^{n/2+1}] + 2^{3/2}(1/2)^{n/2+1} \\ &= 1/3[1 + \sqrt{2}(1/2)^{n/2+1}]. \end{aligned}$$

